Supplemental errata to:

Microbiologic diagnostic tests when asymptomatic carriers are present

Aspects of the use of conventional throat and nasopharyngeal culture as examples by Ronny K Gunnarsson (ISBN 91-628-4746-5)

Another errata (errata_d.pdf) deals with errors in the dissertation. This supplemental errata deals with original publications attached to the printed version of the dissertation.

Publication: Page	Description	At present	Replace with
IV: 2	URL at the bottom of summary	http://www.infovice.se/fou/epv	http://www.infovoice.se/fou/epv
IV: 10	Predictive values at the bottom of the page	91% (74-100%) 96% (88-100%)	93% (78-100%) 95% (87-100%)
IV: 24	In three sentences at the bottom of this page	$1 - \hat{a}$ $1 - 2\hat{a}$ $\hat{a} = \frac{\alpha}{2}$ $1 - \hat{a}$	$1 - \mathbf{b}$ $1 - 2\mathbf{b}$ $\mathbf{b} = \mathbf{a}/2$ $1 - \mathbf{a}$
IV: 25	The first confidence interval in Table 1	(93-99%)	(93-100%)
V:1	Number before Clinical Micro- biology, Central Hospital Växsjö	1	3
V: 2	In Abstract, Objective	eight years	nine years
V: 2	In Abstract, Design	1991-2000	1991-1999
V: 2	In Abstract, Design	11 of 12 2000 laboratories	11 of 12 laboratories
V: 2	In Abstract, Conclusion	stable over time for each area it.	stable over time for each area.
V: 4	On the fifth line	swab samples, the greater the proportion	swab samples one may expect that a greater proportion